

Rational wave solutions to a generalized (2+1)-dimensional Hirota bilinear equation

K. Hosseini^{1*}, M. Mirzazadeh^{2*}, M. Aligoli³, M. Eslami⁴, J.G. Liu⁵

¹Department of Mathematics, Rasht Branch, Islamic Azad University, Rasht, Iran

²Department of Engineering Sciences, Faculty of Technology and Engineering, East of Guilan, University of Guilan, P.C. 44891-63157 Rudsar-Vajargah, Iran

³Department of Applied Mathematics, Faculty of Mathematical Sciences, University of Guilan, P.O. Box 41335-1914, Guilan, Rasht, Iran

⁴Department of Mathematics, Faculty of Mathematical Sciences, University of Mazandaran, Babolsar, Iran

⁵College of Computer, Jiangxi University of Traditional Chinese Medicine, Jiangxi 330004, China

*Corresponding Authors: kamyar_hosseini@yahoo.com, mirzazadehs2@guilan.ac.ir

Abstract: A generalized form of (2+1)-dimensional Hirota bilinear (2D-HB) equation is considered herein in order to study nonlinear waves in fluids and oceans. The present goal is carried out through adopting the simplified Hirota's method as well as ansatz approaches to retrieve a bunch of rational wave structures from multiple soliton solutions to breather, rational, and complexiton solutions. Some figures corresponding to a series of rational wave structures are provided, illustrating the dynamics of the obtained solutions. The results of the present paper indeed help to reveal the existence of rational wave structures of different types for the 2D-HB equation.

Keywords: (2+1)-dimensional Hirota bilinear equation; Generalized form; Simplified Hirota's method; Ansatz approaches; Multiple soliton, breather, rational, and complexiton solutions

1. Introduction

One of the attractive subjects in the areas of mathematical physics is to look for rational wave solutions of differential equations. There are different rational wave solutions to differential equations; for example, multiple soliton solutions, breather solutions, rogue solutions, and so on. During the last decades, many effective methods such as inverse scattering method [1], Hirota's direct method [2], multiple exp-function method [3-6], simplified Hirota's method [7-15], and ansatz methods [16-20] have been exerted to find rational wave solutions of differential equations. More articles can be found in [21-33].

The basic goal of this paper is to study a generalized 2D-HB equation describing nonlinear waves in fluids and oceans as follows [34]

$$\phi_{yt} + c_1(\phi_{xxxxy} + 6\phi_x\phi_y + 3\phi_{xy}\phi + 3\phi_{xx} \int \phi_y dx) + c_2\phi_{yy} = 0, \quad (1)$$

or

$$\phi_{yt} + c_1(\phi_{xxxxy} + 6\phi_x\phi_y + 3\phi_{xy}\phi + 3\phi_{xx}\psi) + c_2\phi_{yy} = 0, \quad \phi_y = \psi_x. \quad (2)$$

which is a generalization of the Hirota bilinear equation studied in [35]. The Hirota bilinear form corresponding to the above nonlinear model (1) is

$$B_{HB}(\varphi) := (D_y D_t + c_1 D_x^3 D_y + c_2 D_y^2) f \cdot f = 2 \left(\varphi \varphi_{yt} - \varphi_y \varphi_t + c_1 (\varphi \varphi_{xxxxy} - 3(\varphi_{xxy} \varphi_x - \varphi_{xy} \varphi_{xx}) - \varphi_y \varphi_{xxx}) + c_2 (\varphi \varphi_{yy} - \varphi_y^2) \right) = 0, \quad (3)$$

under the specific transformation

$$\phi = 2(\ln \varphi)_{xx}.$$

Recently, Hua et al. investigated the interaction behavior associated with the above generalized 2D-HB equation in [34]. They constructed two types of interaction solutions through adopting different test functions as below

$$\varphi = g^2 + h^2 + ke^l + a_{12}, \quad \text{and} \quad \varphi = g^2 + h^2 + \cosh(l) + a_{12}, \quad (4)$$

where

$$g = a_1 x + a_2 y + a_3 t + a_4, \quad h = a_5 x + a_6 y + a_7 t + a_8, \quad l = a_9 x + a_{10} y + a_{11} t. \quad (5)$$

In order to advance the studies on the generalized 2D-HB equation (1); in this paper, the simplified Hirota's method as well as different ansatz approaches are utilized formally to retrieve a bunch of rational wave structures from multiple soliton solutions to breather, rational, and complexiton solutions.

2. The generalized 2D-HB equation and its rational waves solutions

In this section, a number of rational wave structures from multiple soliton solutions to breather, rational, and complexiton solutions for the generalized 2D-HB equation are derived using the simplified Hirota's method as well as ansatz approaches.

2.1. Multiple soliton solutions

In order to procure multiple soliton solutions of the generalized 2D-HB equation, first the expressions

$$\phi = e^{\theta_i}, \quad \theta_i = \kappa_i x + \tau_i y - \zeta_i t, \quad (6)$$

are substituted into the linear terms of the generalized 2D-HB equation; and then, the resulting equation is solved for obtaining the dispersion relation ζ_i . This leads to

$$\zeta_i = c_1 \kappa_i^3 + c_2 \tau_i, \quad (7)$$

and therefore, the phase variables θ_i , $1 \leq i \leq N$ can be written as

$$\theta_i = \kappa_i x + \tau_i y - (c_1 \kappa_i^3 + c_2 \tau_i) t, \quad 1 \leq i \leq N. \quad (8)$$

Now, by inserting the logarithmic transformations

$$\phi = R(\ln \varphi)_{xx}, \quad \psi = R(\ln \varphi)_{xy}, \quad (9)$$

into (2) where the auxiliary function $\varphi(x, y, t)$ is defined as

$$\varphi = 1 + e^{\kappa_1 x + \tau_1 y - (c_1 \kappa_1^3 + c_2 \tau_1) t}, \quad (10)$$

we find that $R = 2$.

Based on this result, a single soliton solution can be obtained as

$$\phi(x, y, t) = \frac{2\kappa_1^2 e^{\kappa_1 x + \tau_1 y - (c_1 \kappa_1^3 + c_2 \tau_1) t}}{1 + e^{\kappa_1 x + \tau_1 y - (c_1 \kappa_1^3 + c_2 \tau_1) t}} - \frac{2\kappa_1^2 e^{2(\kappa_1 x + \tau_1 y - (c_1 \kappa_1^3 + c_2 \tau_1) t)}}{(1 + e^{\kappa_1 x + \tau_1 y - (c_1 \kappa_1^3 + c_2 \tau_1) t})^2}, \quad (11)$$

$$\psi(x, y, t) = \frac{2\kappa_1 \tau_1 e^{\kappa_1 x + \tau_1 y - (c_1 \kappa_1^3 + c_2 \tau_1) t}}{1 + e^{\kappa_1 x + \tau_1 y - (c_1 \kappa_1^3 + c_2 \tau_1) t}} - \frac{2\kappa_1 \tau_1 e^{2(\kappa_1 x + \tau_1 y - (c_1 \kappa_1^3 + c_2 \tau_1) t)}}{(1 + e^{\kappa_1 x + \tau_1 y - (c_1 \kappa_1^3 + c_2 \tau_1) t})^2}. \quad (12)$$

For the double soliton solution, the following auxiliary function is considered

$$\varphi = 1 + e^{\theta_1} + e^{\theta_2} + a_{12} e^{\theta_1 + \theta_2}, \quad (13)$$

in which the phase variables θ_i , $i = 1, 2$ are as before and the phase shift a_{12} is an unknown. After performing some calculations, a double soliton solution is gained as

$$\phi = 2(\ln \varphi)_{xx}, \quad \psi = 2(\ln \varphi)_{xy}, \quad (14)$$

where the phase shift a_{12} is

$$a_{12} = \frac{\kappa_1\tau_1 - \kappa_1\tau_2 - \kappa_2\tau_1 + \kappa_2\tau_2}{\kappa_1\tau_1 + \kappa_1\tau_2 + \kappa_2\tau_1 + \kappa_2\tau_2}. \quad (15)$$

Finally, a triple soliton solution to the generalized 2D-HB equation is derived as

$$\phi = 2(\ln \varphi)_{xx}, \quad \psi = 2(\ln \varphi)_{xy}, \quad (16)$$

in which the auxiliary function, the phase shifts, and the phase variables are defined as

$$\varphi = 1 + e^{\theta_1} + e^{\theta_2} + e^{\theta_3} + a_{12}e^{\theta_1+\theta_2} + a_{13}e^{\theta_1+\theta_3} + a_{23}e^{\theta_2+\theta_3} + a_{12}a_{13}a_{23}e^{\theta_1+\theta_2+\theta_3}, \quad (17)$$

$$a_{ij} = \frac{\kappa_i\tau_i - \kappa_i\tau_j - \kappa_j\tau_i + \kappa_j\tau_j}{\kappa_i\tau_i + \kappa_i\tau_j + \kappa_j\tau_i + \kappa_j\tau_j}, \quad 1 \leq i, j \leq 3, \quad (18)$$

$$\theta_i = \kappa_i x + \tau_i y - (c_1\kappa_i^3 + c_2\tau_i)t, \quad 1 \leq i \leq 3. \quad (19)$$

It is easy to show that the following multiple complex soliton solutions to the generalized 2D-HB equation can be constructed

$$\phi_{1,2,3} = 2(\ln \varphi_{1,2,3})_{xx}, \quad \psi_{1,2,3} = 2(\ln \varphi_{1,2,3})_{xy}, \quad (20)$$

in which

$$\varphi_1 = I + e^{\theta_1}, \quad I = \sqrt{-1}, \quad (21)$$

$$\varphi_2 = I + e^{\theta_1} + e^{\theta_2} - I a_{12} e^{\theta_1+\theta_2}, \quad I = \sqrt{-1}, \quad (22)$$

$$\varphi_3 = I + e^{\theta_1} + e^{\theta_2} + e^{\theta_3} - I a_{12} e^{\theta_1+\theta_2} - I a_{13} e^{\theta_1+\theta_3} - I a_{23} e^{\theta_2+\theta_3} - a_{12} a_{13} a_{23} e^{\theta_1+\theta_2+\theta_3}, \quad I = \sqrt{-1}, \quad (23)$$

and

$$a_{ij} = \frac{\kappa_i\tau_i - \kappa_i\tau_j - \kappa_j\tau_i + \kappa_j\tau_j}{\kappa_i\tau_i + \kappa_i\tau_j + \kappa_j\tau_i + \kappa_j\tau_j}, \quad 1 \leq i, j \leq 3, \quad (24)$$

$$\theta_i = \kappa_i x + \tau_i y - (c_1\kappa_i^3 + c_2\tau_i)t, \quad 1 \leq i \leq 3. \quad (25)$$

The plots of single, double, and triple soliton solutions have been provided in Figures 1-3, illustrating the dynamics of the multiple solutions.

Figure 1 shows a bright soliton wave whereas Figure 2 demonstrates the interaction of two bright soliton waves. Furthermore, the interaction of three bright soliton waves including two strong and one weak waves has been illustrated in Figure 3.




Fig. 1. Single soliton solution when $\kappa_1 = 1$, $\tau_1 = 2$, and $t = 0$.




Fig. 2. Double soliton solution when $\kappa_1 = -3.5$, $\tau_1 = 10$, $\kappa_2 = 3$, $\tau_2 = -1$, and $t = 0$.




Fig. 3. Triple soliton solution when $\kappa_1 = 3$, $\tau_1 = -2$, $\kappa_2 = -5$, $\tau_2 = -9$, $\kappa_3 = -8$, $\tau_3 = 20$, and $t = 0$.

2.2. Breather and rational solutions

In the present subsection, first, the breather solution of the generalized 2D-HB equation is acquired by adopting a specific ansatz approach which is a combination of exponential functions and a trigonometric function as follows

$$\varphi = e^{-kX} + b_0 \cos(hY) + b_1 e^{kX}, \quad (26)$$

in which

$$X = a_1 x + a_2 y + a_3 t + a_4, \quad Y = a_5 x + a_6 y + a_7 t + a_8, \quad (27)$$

and k , b_0 , h , b_1 and a_i , $1 \leq i \leq 8$ are constants to be computed. Inserting Eq. (26) into Eq. (3) and performing some calculations, yields the following nonlinear algebraic system

$$2h^4 a_5^3 a_6 b_0 c_1 - 6h^2 k^2 a_1^2 a_5 a_6 b_0 c_1 - 6h^2 k^2 a_1 a_2 a_5^2 b_0 c_1 + 2k^4 a_1^3 a_2 b_0 c_1 - 2h^2 a_6^2 b_0 c_2 + 2k^2 a_2^2 b_0 c_2 - 2h^2 a_6 a_7 b_0 + 2k^2 a_2 a_3 b_0 = 0, \quad (28)$$

$$2h^4 a_5^3 a_6 b_0 b_1 c_1 - 6h^2 k^2 a_1^2 a_5 a_6 b_0 b_1 c_1 - 6h^2 k^2 a_1 a_2 a_5^2 b_0 b_1 c_1 + 2k^4 a_1^3 a_2 b_0 b_1 c_1 - 2h^2 a_6^2 b_0 b_1 c_2 + 2k^2 a_2^2 b_0 b_1 c_2 - 2h^2 a_6 a_7 b_0 b_1 + 2k^2 a_2 a_3 b_0 b_1 = 0, \quad (29)$$

$$-6h^3 k a_1 a_5^2 a_6 b_0 b_1 c_1 - 2h^3 k a_2 a_5^2 b_0 b_1 c_1 + 2h k^3 a_1^3 a_6 b_0 b_1 c_1 + 6h k^3 a_1^2 a_2 a_5 b_0 b_1 c_1 + 4h k a_2 a_6 b_0 b_1 c_2 + 2h k a_2 a_7 b_0 b_1 + 2h k a_3 a_6 b_0 b_1 = 0, \quad (30)$$

$$6h^3ka_1a_5^2a_6b_0c_1 + 2h^3ka_2a_5^3b_0c_1 - 2hk^3a_1^3a_6b_0c_1 - 6hk^3a_1^2a_2a_5b_0c_1 - 4hka_2a_6b_0c_2 - 2hka_2a_7b_0 - 2hka_3a_6b_0 = 0, \quad (31)$$

$$8h^4a_5^3a_6b_0^2c_1 + 32k^4a_1^3a_2b_1c_1 - 2h^2a_6^2b_0^2c_2 - 2h^2a_6a_7b_0^2 + 8k^2a_2^2b_1c_2 + 8k^2a_2a_3b_1 = 0, \quad (32)$$

whose solution is

$$a_3 = 3h^2a_1a_5^2c_1 - k^2a_1^3c_1 - a_2c_2, \quad a_7 = h^2a_5^3c_1 - 3k^2a_1^2a_5c_1 - a_6c_2, \quad b_1 = -\frac{a_5h^2a_6b_0^2}{4a_1k^2a_2}. \quad (33)$$

Now, a breather solution is obtained as

$$\phi = 2(\ln \varphi)_{xx}, \quad \psi = 2(\ln \varphi)_{xy}, \quad (34)$$

where

$$\varphi = e^{-k(a_1x+a_2y+a_3t+a_4)} + b_0 \cos(h(a_5x + a_6y + a_7t + a_8)) + b_1 e^{k(a_1x+a_2y+a_3t+a_4)}, \quad (35)$$

and a_3 , a_7 , and b_1 have been defined in (33).

The plots of the breather solution (34) have been depicted in Figure 4, demonstrating its dynamical behavior. It is evident that Figure 4 presents the interaction of bright and dark waves with a periodic property.




Fig. 4. Breather solution when $a_1 = 1$, $a_2 = 2$, $a_4 = 0$, $a_5 = -2$, $a_6 = 1$, $a_8 = 0$, $k = 1$, $h = -2$, $b_0 = -1$, and $t = 0$.

Now, the rational solution is obtained from the breather solution (34) in a specific way.

Remark 1. It should be noted that by setting $a_1 = a_5$, $a_2 = -a_6$, $b_0 = -2$, and $h = k$ in (34) and letting $k \rightarrow 0$, a rational solution to the generalized 2D-HB equation can be constructed as

$$\phi(x, y, t) = 8a_5^2 \left(\frac{1}{x^2+y^2} - \frac{(x+y)^2}{(x^2+y^2)^2} \right), \quad (36)$$

$$\psi(x, y, t) = \frac{8a_5 a_6 (x^2 - y^2)}{(x^2 + y^2)^2}, \quad (37)$$

in which $\mathcal{X} = a_5 x - a_6 y + c_2 a_6 t + a_4$ and $\mathcal{Y} = a_5 x + a_6 y - c_2 a_6 t + a_8$.

To show it, by considering $a_1 = a_5$, $a_2 = -a_6$, $b_0 = -2$, and $h = k$; it is obvious that the breather solution (34) can be written as

$$\phi(x, y, t) = \frac{2a_5^2 k^2 (\cosh(kX) + \cos(kY))}{\cosh(kX) - \cos(kY)} - \frac{2(a_5 k (\sinh(kX) + \sin(kY)))^2}{(\cosh(kX) - \cos(kY))^2}, \quad (38)$$

$$\psi(x, y, t) = -2a_5 a_6 k^2 + \frac{2a_5 a_6 k^2 (\sinh^2(kX) - \sin^2(kY))}{(\cosh(kX) - \cos(kY))^2}. \quad (39)$$

Applying the Taylor expansion results in

$$\phi(x, y, t) = \frac{2a_5^2 k^2 \left(\left(1 + \frac{k^2 X^2}{2!} + \dots \right) + \left(1 - \frac{k^2 Y^2}{2!} + \dots \right) \right)}{\left(1 + \frac{k^2 X^2}{2!} + \dots \right) - \left(1 - \frac{k^2 Y^2}{2!} + \dots \right)} - \frac{2 \left(a_5 k \left(\left(kX + \frac{k^3 X^3}{3!} + \dots \right) + \left(kY - \frac{k^3 Y^3}{3!} + \dots \right) \right) \right)^2}{\left(\left(1 + \frac{k^2 X^2}{2!} + \dots \right) - \left(1 - \frac{k^2 Y^2}{2!} + \dots \right) \right)^2}, \quad (40)$$

$$\psi(x, y, t) = -2a_5 a_6 k^2 + \frac{2a_5 a_6 k^2 \left(\left(kX + \frac{k^3 X^3}{3!} + \dots \right)^2 - \left(kY - \frac{k^3 Y^3}{3!} + \dots \right)^2 \right)}{\left(\left(1 + \frac{k^2 X^2}{2!} + \dots \right) - \left(1 - \frac{k^2 Y^2}{2!} + \dots \right) \right)^2}. \quad (41)$$

Now, when k tends to zero; we find a rational solution as

$$\phi(x, y, t) = 8a_5^2 \left(\frac{1}{x^2+y^2} - \frac{(x+y)^2}{(x^2+y^2)^2} \right), \quad (42)$$

$$\psi(x, y, t) = \frac{8a_5 a_6 (x^2 - y^2)}{(x^2 + y^2)^2}, \quad (43)$$

in which $\mathcal{X} = a_5 x - a_6 y + c_2 a_6 t + a_4$ and $\mathcal{Y} = a_5 x + a_6 y - c_2 a_6 t + a_8$.

The above rational solution is plotted graphically in Figure 5 for $a_4 = 0$, $a_5 = -2$, $a_6 = 1$, $a_8 = 0$, and $t = 0$.




Fig. 5. Rational solution when $a_4 = 0$, $a_5 = -2$, $a_6 = 1$, $a_8 = 0$, and $t = 0$.

2.3. Complexiton solutions

This subsection deals with complexiton solutions of the generalized 2D-HB equation. For this purpose, a test function is exerted as

$$\varphi = 1 + 2e^{\theta_1} \cos(\theta_2) + a_{12} e^{2\theta_1}, \quad (44)$$

in which

$$\theta_1 = k_1 x + r_1 y + w_1 t, \quad \theta_2 = k_2 x + r_2 y + w_2 t, \quad (45)$$

and w_1 and w_2 satisfy

$$P(k, r, w) = P(\bar{k}, \bar{r}, \bar{w}) = 0, \quad k = k_1 + I k_2, \quad r = r_1 + I r_2, \quad w = w_1 + I w_2, \quad P = yt + c_1 x^3 y + c_2 y^2, \quad I = \sqrt{-1}. \quad (46)$$

Here, the polynomial P has been defined owing to the Hirota operator (3). It is easy to prove that

$$w_1 = -c_1 k_1^3 + 3c_1 k_1 k_2^2 - c_2 r_1, \quad (47)$$

$$w_2 = c_1 k_2^3 - 3c_1 k_1^2 k_2 - c_2 r_2. \quad (48)$$

The unknown a_{12} also can be gained as

$$a_{12} = -\frac{P(2Ik_2, 2Ir_2, 2Iw_2)}{P(2k_1, 2r_1, 2w_1)} = -\frac{16c_1 k_2^3 r_2 - 4c_2 r_2^2 - 4r_2(-3c_1 k_1^2 k_2 + c_1 k_2^3 - c_2 r_2)}{16c_1 k_1^3 r_1 + 4c_2 r_1^2 + 2r_1(-2c_1 k_1^3 + 6c_1 k_1 k_2^2 - 2c_2 r_1)}. \quad (49)$$

Now, a complexiton solution to the generalized 2D-HB equation is obtained as

$$\phi = 2(\ln \varphi)_{xx}, \quad \psi = 2(\ln \varphi)_{xy}, \quad (50)$$

in which

$$\varphi = 1 + 2e^{\theta_1} \cos(\theta_2) + a_{12} e^{2\theta_1}, \quad (51)$$

$$\theta_1 = k_1 x + r_1 y + (-c_1 k_1^3 + 3c_1 k_1 k_2^2 - c_2 r_1) t, \quad \theta_2 = k_2 x + r_2 y + (c_1 k_2^3 - 3c_1 k_1^2 k_2 - c_2 r_2) t, \quad (52)$$

$$a_{12} = -\frac{16c_1 k_2^3 r_2 - 4c_2 r_2^2 - 4r_2(-3c_1 k_1^2 k_2 + c_1 k_2^3 - c_2 r_2)}{16c_1 k_1^3 r_1 + 4c_2 r_1^2 + 2r_1(-2c_1 k_1^3 + 6c_1 k_1 k_2^2 - 2c_2 r_1)}. \quad (53)$$

It is worth mentioning that the following complex complexiton solution to the generalized 2D-HB equation can be extracted

$$\phi = 2(\ln \varphi)_{xx}, \quad \psi = 2(\ln \varphi)_{xy}, \quad (54)$$

in which

$$\varphi = I + 2e^{\theta_1} \cos(\theta_2) - I a_{12} e^{2\theta_1}, \quad I = \sqrt{-1}, \quad (55)$$

$$\theta_1 = k_1 x + r_1 y + (-c_1 k_1^3 + 3c_1 k_1 k_2^2 - c_2 r_1) t, \quad \theta_2 = k_2 x + r_2 y + (c_1 k_2^3 - 3c_1 k_1^2 k_2 - c_2 r_2) t, \quad (56)$$

$$a_{12} = -\frac{16c_1 k_2^3 r_2 - 4c_2 r_2^2 - 4r_2(-3c_1 k_1^2 k_2 + c_1 k_2^3 - c_2 r_2)}{16c_1 k_1^3 r_1 + 4c_2 r_1^2 + 2r_1(-2c_1 k_1^3 + 6c_1 k_1 k_2^2 - 2c_2 r_1)}. \quad (57)$$

The above complexiton solution has been portrayed graphically in Figure 6 for suitable parameters.

2.4. Hyperbolic solutions

In this subsection, hyperbolic solutions of the generalized 2D-HB equation are obtained with the use of ansatz methods. To start, let's consider the solution of the generalized 2D-HB equation as

$$\phi = A_0 + A_1 \tanh(\kappa x + \tau y + \omega t)^2, \quad (58)$$

$$\psi = \frac{\tau}{\kappa} (A_0 + A_1 \tanh(\kappa x + \tau y + \omega t)^2). \quad (59)$$

By setting Eqs. (58) and (59) in the system (2) and performing some calculations, we find

$$A_1 = -2\kappa^2, \quad \omega = 8c_1 \kappa^3 - 6A_0 c_1 \kappa - c_2 \tau. \quad (60)$$




Fig. 6. Complexiton solution when $k_1 = -2$, $k_2 = -2$, $r_1 = -2$, $r_2 = 10$, and $t = 0$.

Therefore, the following hyperbolic solution to the generalized 2D-HB equation is obtained

$$\phi = A_0 - 2\kappa^2 \tanh(\kappa x + \tau y + (8c_1\kappa^3 - 6A_0c_1\kappa - c_2\tau)t)^2, \quad (61)$$

$$\psi = \frac{\tau}{\kappa} (A_0 - 2\kappa^2 \tanh(\kappa x + \tau y + (8c_1\kappa^3 - 6A_0c_1\kappa - c_2\tau)t)^2). \quad (62)$$

It can be readily shown that another hyperbolic solution can be derived as

$$\phi = A_0 - 2\kappa^2 \coth(\kappa x + \tau y + (8c_1\kappa^3 - 6A_0c_1\kappa - c_2\tau)t)^2, \quad (63)$$

$$\psi = \frac{\tau}{\kappa} (A_0 - 2\kappa^2 \coth(\kappa x + \tau y + (8c_1\kappa^3 - 6A_0c_1\kappa - c_2\tau)t)^2). \quad (64)$$

Now, let us consider the following hyperbolic solution to the generalized 2D-HB equation

$$\phi = A_0 + A_1 \operatorname{sech}(\kappa x + \tau y + \omega t)^2, \quad (65)$$

$$\psi = \frac{\tau}{\kappa} (A_0 + A_1 \operatorname{sech}(\kappa x + \tau y + \omega t)^2). \quad (66)$$

Substituting Eqs. (65) and (66) into the system (2) and performing some calculations, results in

$$A_1 = 2\kappa^2, \quad \omega = -4c_1\kappa^3 - 6A_0c_1\kappa - c_2\tau. \quad (67)$$

Therefore, the following hyperbolic solution to the generalized 2D-HB equation is derived

$$\phi = A_0 + 2\kappa^2 \operatorname{sech}(\kappa x + \tau y + (-4c_1\kappa^3 - 6A_0c_1\kappa - c_2\tau)t)^2, \quad (68)$$

$$\psi = \frac{\tau}{\kappa} (A_0 + 2\kappa^2 \operatorname{sech}(\kappa x + \tau y + (-4c_1\kappa^3 - 6A_0c_1\kappa - c_2\tau)t)^2). \quad (69)$$

In a parallel manner, it can be demonstrated that

$$\phi = A_0 - 2\kappa^2 \operatorname{csch}(\kappa x + \tau y + (-4c_1\kappa^3 - 6A_0c_1\kappa - c_2\tau)t)^2, \quad (70)$$

$$\psi = \frac{\tau}{\kappa} (A_0 - 2\kappa^2 \operatorname{csch}(\kappa x + \tau y + (-4c_1\kappa^3 - 6A_0c_1\kappa - c_2\tau)t)^2), \quad (71)$$

is the solution of the generalized 2D-HB equation.

The plots of the hyperbolic solution (68) have been provided in Figure 7. Obviously, Figure 7 shows a bright soliton wave solution.




Fig. 7. Hyperbolic solution (68) when $A_0 = 1$, $\kappa = 1$, $\tau = 1$, and $t = 0$.

Remark 2. Different classes of rational wave solutions have been reported in the present paper which definitely are different from the exact solutions of Ref. [34].

3. Concluding remarks

A generalized (2+1)-dimensional Hirota bilinear equation which describes nonlinear waves in fluids and oceans was comprehensively explored, in this paper. The current goal was performed by exerting the simplified Hirota's method and ansatz approaches as reliable techniques. Different classes of rational wave

structures such as multiple soliton solutions, breather solutions, rational solutions, and complexiton solutions to the generalized 2D-HB equation were successfully obtained. Physical behaviors of a series of rational wave structures were demonstrated graphically. The results of the current paper certainly helped to reveal the existence of rational wave structures of different types for the 2D-HB equation.

References

1. C.S. Gardner, J.M. Greene, M.D. Kruskal, R.M. Miura, Method for solving the Korteweg-deVries equation, *Physical Review Letters* 19 (1967) 1095–1097.
2. R. Hirota, *The direct method in soliton theory*, Cambridge University Press, Cambridge, 2004.
3. W.X. Ma, T. Huang, Y. Zhang, A multiple exp-function method for nonlinear differential equations and its application, *Physica Scripta* 82 (2010) 065003.
4. W.X. Ma, Z. Zhu, Solving the (3+1)-dimensional generalized KP and BKP equations by the multiple exp-function algorithm, *Applied Mathematics and Computation* 218 (2012) 11871–11879.
5. A.R. Adem, M. Mirzazadeh, Q. Zhou, K. Hosseini, Multiple soliton solutions of the Sawada–Kotera equation with a nonvanishing boundary condition and the perturbed Korteweg de Vries equation by using the multiple exp-function scheme, *Advances in Mathematical Physics* 2019 (2019) 3175213.
6. M. Inc, K. Hosseini, M. Samavat, M. Mirzazadeh, M. Eslami, M. Moradi, D. Baleanu, N -wave and other solutions to the B-type Kadomtsev–Petviashvili equation, *Thermal Science* 23 (2019) 2027–2035.
7. W. Hereman, W. Zhuang, Symbolic software for soliton theory, *Acta Applicandae Mathematicae* 39 (1995) 361–378.
8. W. Hereman, A. Nuseir, Symbolic methods to construct exact solutions of nonlinear partial differential equations, *Mathematics and Computers in Simulation* 43 (1997) 13–27.
9. A.M. Wazwaz, Multiple-soliton solutions for the generalized (1+1)-dimensional and the generalized (2+1)-dimensional Ito equations, *Applied Mathematics and Computation* 202 (2008) 840–849.
10. A.M. Wazwaz, L. Kaur, New integrable Boussinesq equations of distinct dimensions with diverse variety of soliton solutions, *Nonlinear Dynamics* 97 (2019) 83–94.
11. A.M. Wazwaz, Multiple complex soliton solutions for the integrable KdV, fifth-order Lax, modified KdV, Burgers, and Sharma–Tasso–Olver equations, *Chinese Journal of Physics* 59 (2019) 372–378.
12. A.M. Wazwaz, Multiple complex and multiple real soliton solutions for the integrable sine-Gordon equation, *Optik* 172 (2018) 622–627.

13. A.M. Wazwaz, Multiple complex soliton solutions for integrable negative-order KdV and integrable negative-order modified KdV equations, *Applied Mathematics Letters* 88 (2019) 1–7.
14. A.M. Wazwaz, Multiple-soliton solutions for extended shallow water wave equations, *Studies in Mathematical Sciences* 1 (2010) 21–29.
15. K. Hosseini, M. Aligoli, M. Mirzazadeh, M. Eslami, J.F. Gómez-Aguilar, Dynamics of rational solutions in a new generalized Kadomtsev–Petviashvili equation, *Modern Physics Letters B* 33 (2019) 1950437.
16. X.X. Du, B. Tian, Y. Yin, Lump, mixed lump-kink, breather and rogue waves for a B-type Kadomtsev–Petviashvili equation, *Waves in Random and Complex Media* (2019), doi. 10.1080/17455030.2019.1566681.
17. W. Tan, H. Dai, Z. Dai, W. Zhong, Emergence and space-time structure of lump solution to the (2+1)-dimensional generalized KP equation, *Pramana Journal of Physics* 89 (2017) 77.
18. C.C. Hu, B. Tian, H.M. Yin, C.R. Zhang, Z. Zhang, Dark breather waves, dark lump waves and lump wave-soliton interactions for a (3+1)-dimensional generalized Kadomtsev–Petviashvili equation in a fluid, *Computers and Mathematics with Applications* 78 (2019) 166–177.
19. R. Pouyanmehr, K. Hosseini, R. Ansari, S.H. Alavi, Different wave structures to the (2+1)-dimensional generalized Bogoyavlensky–Konopelchenko equation, *International Journal of Applied and Computational Mathematics* 5 (2019) 149.
20. A.M. Wazwaz, Two wave mode higher-order modified KdV equations: Essential conditions for multiple soliton solutions to exist, *International Journal of Numerical Methods for Heat & Fluid Flow* 27 (2017) 2223–2230.
21. Y. Zhou, W.X. Ma, Complexiton solutions to soliton equations by the Hirota method, *Journal of Mathematical Physics* 58 (2017) 101511.
22. J.G. Liu, J.Q. Du, Z.F. Zeng, B. Nie, New three-wave solutions for the (3+1)-dimensional Boiti–Leon–Manna–Pempinelli equation, *Nonlinear Dynamics* 88 (2017) 655–661.
23. J.G. Liu, M. Eslami, H. Rezazadeh, M. Mirzazadeh, Rational solutions and lump solutions to a non-isospectral and generalized variable-coefficient Kadomtsev–Petviashvili equation, *Nonlinear Dynamics* 95 (2019) 1027–1033.
24. J.G. Liu, Q. Ye, Stripe solitons and lump solutions for a generalized Kadomtsev–Petviashvili equation with variable coefficients in fluid mechanics, *Nonlinear Dynamics* 96 (2019) 23–29.

25. J.G. Liu, Y. He, Abundant lump and lump-kink solutions for the new (3+1)-dimensional generalized Kadomtsev–Petviashvili equation, *Nonlinear Dynamics* 92 (2018) 1103–1108.
26. R. Zhang, S. Bilige, T. Fang, T. Chaolu, New periodic wave, cross-kink wave and the interaction phenomenon for the Jimbo–Miwa-like equation, *Computers and Mathematics with Applications* 78 (2019) 754–764.
27. L.L. Feng, T.T. Zhang, Breather wave, rogue wave and solitary wave solutions of a coupled nonlinear Schrödinger equation, *Applied Mathematics Letters* 78 (2018) 133–140.
28. J. He, Y. Li, Designable integrability of the variable coefficient nonlinear Schrödinger equations, *Studies in Applied Mathematics* 126 (2011) 1–15.
29. J. He, L. Wang, L. Li, K. Porsezian, R. Erdlyi, Few-cycle optical rogue waves: Complex modified Korteweg-de Vries equation, *Physical Review E* 89 (2014) 062917.
30. J.S. He, E.G. Charalampidis, P.G. Kevrekidis, D.J. Frantzeskakis, Rogue waves in nonlinear Schrödinger models with variable coefficients: Application to Bose–Einstein condensates, *Physics Letters A* 378 (2014) 577–583.
31. J. He, S. Xu, Y. Cheng, The rational solutions of the mixed nonlinear Schrödinger equation, *AIP Advances* 5 (2015) 017105.
32. M.S. Osman, M. Inc, J.G. Liu, K. Hosseini, A. Yusuf, Different wave structures and stability analysis for the generalized (2+1)-dimensional Camassa–Holm–Kadomtsev–Petviashvili equation, *Physica Scripta* 95 (2020) 035229.
33. M.S. Osman, A.M. Wazwaz, A general bilinear form to generate different wave structures of solitons for a (3+1)-dimensional Boiti–Leon–Manna–Pempinelli equation, *Mathematical Methods in the Applied Sciences* 42 (2019) 6277–6283.
34. Y.F. Hua, B.L. Guo, W.X. Ma, X. Lü, Interaction behavior associated with a generalized (2+1)-dimensional Hirota bilinear equation for nonlinear waves, *Applied Mathematical Modelling* 74 (2019) 184–198.
35. X. Lü, W.X. Ma, Study of lump dynamics based on a dimensionally reduced Hirota bilinear equation, *Nonlinear Dynamics* 85 (2016) 1217–1222.